

**OSELTAMIVIR TREATMENT
Influenza A and B Infection
Residential Care**

 Weight (kg)

--

 Bulleted orders are initiated by default, unless crossed out and initialed by the physician/prescriber. Boxed orders () require physician/prescriber check mark () to be initiated.

1. LABORATORY
NOTE: The estimated GFR (eGFR) is **AUTOMATICALLY REPORTED** for each creatinine

- Creatinine retesting not required if (*check all that apply*):
 - Estimated GFR greater than 60 mL/minute (in last 12 months) and no renal dysfunction suspected by physician
 - Recent creatinine available (previous 4 weeks)
 - Chronic hemodialysis or peritoneal dialysis patient
- Obtain a creatinine if (*check all that apply*):
 - No creatinine tested in the last 12 months
 - Estimated GFR less than 60 mL/minute OR renal dysfunction suspected by physician
- Estimated GFR: _____ mL/minute Date: _____ (DD/MM/YY)

2. MEDICATIONS

- Do not administer oseltamivir: Indicate reason(s) (*check all that apply*):
 - Oseltamivir allergy or severe intolerance
 - Patient, caregiver or substitute decision-maker refusal
 - Physician decision
- TREATMENT (*for symptomatic residents only*)
 - Initiate oseltamivir TREATMENT within 48 hours of symptom onset

Select one	Estimated GFR (mL/min.)	Adult oseltamivir TREATMENT [13 yrs and older]
<input type="checkbox"/>	Greater than 60	75 mg PO BID × 5 days
<input type="checkbox"/>	31–60	30 mg PO BID × 5 days
<input type="checkbox"/>	10–30	30 mg PO DAILY × 5 days
<input type="checkbox"/>	Peritoneal Dialysis	30 mg PO × 1 dose only (give before dialysis)
<input type="checkbox"/>	Hemodialysis	30 mg PO BEFORE DIALYSIS, then 30 mg AFTER EACH DIALYSIS SESSION × 5 consecutive days

Date (dd/mm/yyyy)	Time	Physician Signature	Printed Name or College ID#
/ /			

Scan or Fax page to Pharmacy