





Kamloops

DOWN contains medetomidine and fluorofentanyl

Looks Like: Purple chunks and pebbles

Sold as: Down, Dope, Fentanyl

Contains: Fluorofentanyl (31%)


Medetomidine (3%)

Risk: High risk of overdose

Dizziness and confusion

Nausea, amnesia, sleepiness, nodding out for a long time

Smoking is not safer than injecting



Medetomidine can lead to complicated and long lasting overdose symptoms. Medetomidine suppresses respiration and has potentially dangerous cardiovascular effects.

The combination of fluorofentanyl and medetomidine can lead to significant sedation. Overdose symptoms may last beyond administration of naloxone.



If you have a drug that looks like this, get it tested before using. Drug checking is available in Kamloops Monday to Friday from 8:30-12:00 and 1:00-4:30 at Ask Wellness- 433 Tranquille Road (back entrance)



Consider these additional tips if you are going to use drugs

- Find more drug checking locations at www.drugchecking.ca
- Be aware of risks if mixing with other drugs, including alcohol
- Use with others around or at an Overdose Prevention Site
- · Start with a small amount and wait before taking more
- Space out your doses
- Carry naloxone and know how to use it
- Call 211 or visit bc211.ca to find services near you
- Get the LifeGuard App lifeguarddh.com

Alert in effect until October 8th 2025

Harm Reduction | Population Health

Local
Substance Use Services
310-MHSU
Kamloops Drug Checking
250-299-7219
Kamloops OPS
250-377-6500

