



Management of Patients Requiring Airborne Isolation IN THE ABSENCE of Airborne Infection Isolation Rooms (AIIRs)

Apply a medical mask to the patient as tolerated.

Apply appropriate Additional Precaution sign(s) as required for the airborne infectious disease (e.g., Airborne & Droplet & Contact Precautions), refer to Syndromic Surveillance Screening toolkit.

Health care providers require fit-tested N95 Respirator and other PPE based on other Additional Precautions and Point of Care Risk Assessment (PCRA).





discharge for 2 hours or per the air clearance or settle time.

Keep Additional Precautions sign in place until settle time has lapsed. Environmental Services to remove the sign after Additional Precautions Discharge Cleaning and Disinfection completed.

Persons needing to enter the room before air clearance time has passed must wear fit-tested N95 respirator.

Contact IPAC to discuss if transfer to facility with available AIIR is recommended. Keep patient in dedicated space until transport. Patient to wear a medical mask for transport as

On transfer: keep room vacant with door closed post- discharge for 2 hours or per the air clearance or settle time.

Keep Additional Precaution sign(s) in place until settle time has lapsed. Environmental Services to remove the sign after Additional Precautions Discharge Cleaning and Disinfection completed.

Persons needing to enter the room before air clearance time has passed must wear fit-tested N95 respirator.

Effective Date	July 2024		
Last Reviewed			
Approved By	IPAC		
Owner	Infection Prevention and Control		
	Date	Section	Revision